



www.jahm.in  
(eISSN-2321-1563)



## ORIGINAL RESEARCH ARTICLE

# CLINICAL STUDY TO EVALUATE THE EFFICACY AND SAFETY OF “HYPERCUM” IN HYPERTENSION AND ASSOCIATED QUALITY OF LIFE - AN OPEN LABELED PROOF OF CONCEPT STUDY

<sup>1</sup>HARISHA. S, <sup>2</sup>KIRAN.VUPPALA, <sup>3</sup>ANIL KUMAR SHARMA

<sup>1,2</sup> ICBio Clinical Research Pvt, Vidyaranyapura, Bangalore - 97, India.

<sup>3</sup> Natural Solutions, #26, MHADA Complex off Link Road, Oshiwara, andheri (w), Mumbai-400053.

Corresponding author Email address: clinical.mgt@icbiocro.com

Access this article online: www.jahm.in

Published by Atreya Ayurveda Publications, Ilkal-587125 (India) all rights reserved.

Received on: 11/04/14, Revised on: 14/04/14, Accepted on: 25/04/14

## ABSTRACT

**Background:** High blood pressure has emerged as a leading cause for death and disability worldwide. Management of hypertension effectively has been a distant reality. Hypercum, a rationale combination of herbs has shown to reduce BP effectively along with other benefits as studied in the ancient literatures and research publication. **Objectives:** The aim of the study is to evaluate the safety and efficacy of “Hypercum” on mild to moderate hypertension, associated with co-morbid conditions. **Methods:** An open labeled nonrandomized proof of concept study enrolled 16 hypertensive patients, who met the selection criteria. The efficacy was assessed by measuring the blood pressure at baseline and on every 15 days till the BP got controlled and then every month and followed up for 84 days. The effect of Hypercum on other co-morbid conditions was also evaluated. The quality of life was also assessed using SF-30. Planned student ‘t’ test was applied. **Results:** At screening (before the enrollment for the study) the mean systolic blood pressure (SBP) was 161.25 mmHg and their mean diastolic blood pressure (DBP) was 106.88mmHg. The mean baseline systolic blood pressure, the mean diastolic blood pressure and the mean arterial blood pressure (MAP) were 152.50 mmHg, 99 mmHg and 113.88 mmHg respectively. The mean value of these parameters SBP, DBP and MAP decreased to 122.50 mmHg, 81.67 mmHg and 92.58 mmHg respectively at visit 5 that is at Day 84. There was significant reduction in systolic blood pressure, diastolic blood pressure and mean arterial blood pressure from baseline to the entire visit that is Day14, Day28, Day56 and Day84. **Conclusion:** “Hypercum” is effective in treating the mild to moderate hypertension associated with co-morbid conditions with positive outcome on the quality of life.

**Key words:** Hypercum, Hypertension, Blood pressure, co-morbid conditions.

## INTRODUCTION

High blood pressure is the number one risk factor for death and disability worldwide<sup>1</sup>. Recent reports indicate that nearly 1 billion adults (more than a quarter of the world’s population) had hypertension in 2000, and this is predicted to increase to 1.56 billion by 2025. Approximately 30% of the population has hypertension, and the prevalence is further increasing. The risk has increased in individuals with co-

morbidities such as diabetes, chronic kidney disease, and coronary artery disease<sup>2</sup>.

Poorly controlled hypertension causes cardiovascular disease, resulting in increased risk of stroke, heart disease (including myocardial infarction, heart failure, and arrhythmias) and kidney disease. Scientific literature available on herbal medicines has revealed that herbs are effective in the cardiovascular system both in terms of efficacy and safety<sup>3</sup>.

Despite evidence of efficacy of antihypertensive agents in treating hypertensive patients, the achievement and maintenance of the target BP goals, still remains a challenge for the treating physicians, with limited armamentarium.

According to present scenario, dependence on natural products is gaining popularity day- by-day to combat various physiological threats including cardiovascular complexities. The use of traditional remedies may be encountered more frequently due to an array of scientific evidence in it's favor<sup>4</sup>. Reports indicate that about 15–20% of individuals on prescription medications also use herbal supplements<sup>5</sup>. Hence it is imperative to promote credible research on the safety and efficacy of herbal treatment for variety of ailments including cardiovascular diseases.

**Hypercum** is a herbal drug which comprises of *Rauvolfia serpentina*, *Nardostachys jatamanshi*, *Allium sativum*, *Ocimum sanctum*, *Curcumin longa*, *Embilica officinalis*, *Azadirachta indica*, *Trigonella foenum graecum*, *Zingiber officinale*, *Tribulus terrestris*, *Withania somnifera*, *Pueraria tuberosa*, *Terminalia arjuna*, *Centella asiatica*, *Mentha arvensis* and *Marsh mint*.

Indian literature has reviewed the efficacy and safety of Hypercum ingredients at various concentrations in many clinical conditions. However, till date, there is lack of evidence on the polyherbal combination 'Hypercum'.

## OBJECTIVES:

**Primary objective:** To evaluate the anti hypertensive efficacy and safety of Hypercum on mild to moderate hypertension

**Secondary objective:** To evaluate the benefits of Hypercum on the co-morbid conditions and its impact on quality of life.

## MATERIALS AND METHODS:

Patients both male and female aged 18-60 years and those willing to give written informed consent were selected. They were with mild-to-moderate essential hypertension.

### Inclusion criteria:

Patients taking anti-hypertensive drugs or patients having known to be a hypertensive, diabetic patients or patients running with renal failure or patients having

metabolic syndrome, female subjects who confirmed non-pregnant status and agreed to comply with proper contraception throughout the study duration and patients willing and able to comply with all trial requirements were included in the study. The subjects who were enrolled in the study were hypertensive from several years and were on treatment therapy to control the blood pressure. The medical history of the 16 patients reported that the mean systolic pressure of 161.25 mmHg and their mean diastolic blood pressure of 106.88mmHg.

### Exclusion criteria:

Patients with severe essential hypertension, significant renal insufficiency, history of cerebro-vascular disease, HIV infection, AIDS, hepatitis B or C, or other immunosuppressive disorders, drug abuse within past 2 years, pregnancy and breast feeding women were excluded from the study.

Study was conducted by non- randomized, open labeled, interventional study by ICBio clinical research. It involved the clinical attendance of the subjects on recruitment and on follow up. Subjects enrolled in the study received study drug (3 gms of Hypercum along with ½ tsp of water after food at bed time) during each visit. Study drug would be the alternative or in addition to the therapies already being used.

The safety and efficacy parameters were compared with baseline and follow-up data with laboratory investigations, demographics and blood pressure which included systolic blood pressure, diastolic blood pressure and also the mean arterial blood pressure were analyzed in the study. Adverse events / side effects were noted for each follow up visit.

## Study outcomes

### Primary outcomes

- Reduction in Systolic blood pressure and Diastolic blood pressure from baseline
- Change in mean arterial blood pressure
- Adverse events resulted in therapy
- Laboratory tests (Haematology, Biochemistry and urine analysis) at baseline and post study

**Secondary outcomes:** Quality of life SF-36 questionnaire

**Visit details:** The patients were screened and enrolled. The enrollment day was considered as the baseline data

and the patient were asked to visit on: Day14, Day28, Day56 and Day84.

**Statistical analysis:** Data analysis was carried out using Statistical Analysis System. Student ‘t’ test for independent samples was used to compare group mean baseline values and response differences (outcomes minus baselines) between the groups. Planned student ‘t’ test for paired values was used to compare outcome versus baseline values with in groups. Significant differences between mean data were determined using  $P < 0.05$ . Quality of life evaluation was done through Chi square test.

**RESULTS:**

**Demographic & other baseline characteristics:** In the study around 18 patients were screened and out of them 16 patients were selected. The other 2 patients were considered as screen failure as they did not meet the inclusion criteria. The enrolled subjects consisted of 8 Males and 8 females (Table 1).

Smoking status			
Never Smoked	Still Smoking	Quit smoking	
12	04	00	
Weight			
Mean Weight of Men		Mean Weight of Women	
77.29 Kg		69.53 kg	
Height			
Mean Height of Men		Mean height of Women	
168.75 cm		157.5 cm	
Family history			
Mother	Father	Both	Sibling
06	01	03	04
Medical history			
Hypertensive subjects	Diabetic subjects	Hypertensive & Diabetic subject	Other Medical History
05	02	01	01 (Hypothyroid)
Surgical history			
Number	Reason for Surgical History		
03	Hysterectomy :01 Renal Calculi : 01 Fractured Right Leg :01		


**Concomitant medication taken by the subject:** Hypercum is a herbal drug which can be taken along with other hypertensive drug without altering the routine blood parameters (Table 2).

**Table 2: Concomitant medication taken by the subject**

Concomitant Medication	No. of patients	Medication intake
<b>Antihypertensive drugs</b> (Telmisartan, Olmesartan, Losartan and Amlodpine)	<b>10</b>	Ongoing
<b>Antidiabetic drugs</b> (Human insulin, Metformin & Glibenclamide)	<b>4</b>	Ongoing


**Efficacy Analyses:** The primary efficacy analyses included systolic blood pressure, diastolic blood pressure and mean arterial blood pressure. These parameters were assessed at 5 visits. The subjects were followed from the baseline to Day 14 (visit 2), Day 28 (Visit 3), Day 56 (Visit 4) and Day 84 (Visit 5) (Figure 1).

**Figure 1: Percentage change in systolic blood pressure**



A significant drop in systolic blood pressure from the baseline was noticed by the end of the trial (Day 84).

**Figure 2: Percentage change in diastolic blood pressure**



A significant drop in diastolic blood pressure from the baseline was noticed by day 84.

**Table 3: Mean arterial blood pressure from baseline to evaluation period**

Visit Day	Mean Value	Change from Base line	% Change from Base line
Base Line	113.88 ± 19.7		
Day 14 (Visit 2)	100.12 ± 9.61	-13.75 ± 15.07	0.12
Day 28 (Visit 3)	94.5 ± 8.06	-19.38 ± 14.49	-5.50
Day 56 (Visit 4)	92.93 ± 6.14	-20.94 ± 15.28	-7.06
Day 84 (Visit 5)	92.18 ± 5.54	-21.69 ± 15.98	-7.81

By Day 84, there was significant drop in mean arterial blood pressure from the baseline.

**Safety analyses:** Safety analysis was carried out through the study and Hypercum was well-tolerated. There were no significant findings found for the study drug. No serious adverse events were reported till date. All laboratory parameters were found normal values at screening visit for all subjects. There were no significant findings at baseline and after the study in any of the patients.

**Benefits on Co-morbid conditions:** Hypertension is not only a single disease but it also consists of various other illness associated with it like head ache, body pain, muscle tenderness, depriving of sound sleep,

breathlessness, stomach discomfort (associated more with other allopathic medication), anxiety and palpitation, mood swings, constipation, chest heaviness.


To assess this illness in the hypertensive subjects, set of questionnaire was prepared which was asked to the subjects at the time of enrollment and subsequently at each visit by the investigator. The result obtained from the questionnaire was then analyzed and it was noticed that at the base line i.e., at enrollment few of the subjects complained about mild body pain, mild breathlessness, mild joint pain, moderate mood swing, mild constipation further when these patients took Hypercum and were followed up later almost all the subjects were cured of the above said illness. (Table 4)

**Table 4: Symptoms captured at screening as well as at final visit**

Symptoms	Screening	Visit 5 (Final Visit)
Lower Back Pain	3	1
Breathlessness	3	0
Constipation	3	0
Mood Swing	2	0
Joint Pain	2	1
Abdominal Discomfort	0	1

**Impact on Quality of life:** The mean values for Physical Function (PF), Role-Physical (RP), Bodily Pain (BP), General Health (GH), Vitality (VT), Social Functioning (SF), Role Emotional (RE) and mental health (MH) have shown improvement after the treatment when compared from baseline. The Physical Component Score (PCS) and Mental Component Score have shown improvement in the quality of life when compared from baseline. The mean values are as shown in Fig 3.

**Fig 3: SF 36 individual function assessment**



**Discussion:** During baseline the subjects were screened according to the inclusion- exclusion criteria's and their blood pressure along with mean arterial blood pressure was taken down. The baseline blood pressure prior to administering the drug was taken and at that point the systolic blood pressure was  $152.50 \pm 12.38$  mmHg, diastolic blood pressure was  $99 \pm 10.03$  mmHg and the mean arterial blood pressure was  $113.88 \pm 19.37$  mm Hg. The systolic pressure after 30 min was  $145.88 \pm 9.56$ , diastolic pressure after 30 min was  $94.13 \pm 9.02$  and the mean arterial blood pressure recorded was  $110.21 \pm 20.30$  mmHg.

The patients were then asked to follow the drug schedule and to come for follow up regularly on Day 14 (Visit 2), Day 28 (Visit 3), Day 56 (visit 4) and Day 84 (Visit 5) from the baseline visit. During these visits the subject's vitals and other physical examinations were conducted. During the last visit that is at Day 84(Visit 5) the subject's blood pressure and the mean arterial blood pressure showed a significant change from the baseline. The systolic blood pressure at Day Finally, it can be concluded from the above study that Hypercum not only is an effective against hypertension and also is safe to be taken along with other prescribed medication which helps in reducing other illness associated with hypertension.

**Conclusion:** Hypercum, a polyherbal combination is effective in achieving the normal blood pressure when administered with other antihypertensive drugs without any changes in blood parameters. Hypercum was also effective in improving the quality of life of hypertensive patients. Hypercum is safe and effective in treating mild to moderate hypertension with added effect on the associated co-morbid conditions.

## References

1. Michael BH, Edgar AJ. Optimizing combination therapy in the management of hypertension: the role of the aliskiren,

84 (Visit 5) was  $122.63 \pm 9.29$  mmHg, the diastolic blood pressure was  $81.75 \pm 5.05$  mmHg and the mean arterial blood pressure from the left ankle was  $92.19 \pm 5.54$  mm Hg. The reduction in the blood pressure and the mean arterial blood pressure from the left ankle showed significant changes.

Further, when these changes were compared with each visit statistically the p value was  $< 0.005$ ; hence we can conclude that Hypercum is a potent polyherbal combination in controlling hypertension.

Hypercum can be taken safely along with other hypertensive drug. Hypercum was also analyzed with respect to illness associated along with hypertension. It was observed that few subjects during screening, subjects complained about various other illness associated with hypertension such as body pain, back pain, constipation, mood swing. These illnesses were noted at the baseline by the investigator and were followed up till the final visit it was noticed that almost all the subjects were cured of this illness.

- amlodipine, and hydrochlorothiazide fixed combination. *Integrated Blood Pressure Control*.2013;6 59–67.
2. Epidemiology of Hypertension. *SUPPLEMENT TO JAPI* .2013 (61):12-13.
3. Herbal Medicine for the Treatment of Cardiovascular Disease. *Arch Intern Med*.1998; 158 (20):2225-2234.
4. Blumenthal M. Herb market levels after five years of boom. *Herbal Gram*.1999; 47:64-65.
5. Kaufman DW, Kelly JP, Rosenberg L, Anderson TE, Mitchell AA. Recent patterns of medication use in ambulatory adult population.

**Cite this article as:** Harisha. S, Kiran.Vuppala, Anil K Sharma, Clinical trial of poly herbal product in the treatment of hypertension. *Journal of Ayurveda and Holistic Medicine (JAHM)*. 2014; 2(4).p.12-16.

Source of support: Nil, Conflict of interest: None Declared.